

Lepton Flavor Violating $\tau \rightarrow 3\mu$ Decay in Type-III Two Higgs Doublet Model

Akihiro Matsuzaki¹ and Hidekazu Tanaka²

*Department of Physics, Rikkyo University,
Nishi-ikebukuro, Toshima-ku Tokyo, Japan, 171*

Abstract

We study the current structure of the lepton flavor violating $\tau \rightarrow 3\mu$ decay in Type-III 2HDM. This model has many coupling constants which affect this decay. We find that each coupling constant corresponds to the different final-state momenta distribution and vice versa. Using this fact, we suggest how to determine the current structure. We also find the upper limit $|\eta_{23}^E \eta_{22}^E| < 0.00022$ in the case that all Higgs bosons except for the lighter CP even neutral one h^0 are decoupled, $M_{h^0} = 115\text{GeV}$ and $\cos \beta = 1/\sqrt{2}$. The observable difference between the MSSM and type-III 2HDM is also discussed.

¹akihiro@rikkyo.ac.jp

²tanakah@rikkyo.ac.jp

1 Introduction

In the Standard Model (SM), which is supported by many experimental data, only the Higgs boson is undiscovered. The Large Hadron Collider (LHC) will start and search it [1]. In LHC, we hope to discover many new particles since they are very important hints to beyond the SM. However these particles may be too heavy to discover in LHC. Even if so, the Higgs boson mass has the upper limit from the unitarity [2]. In the Higgs sector, many models are suggested e.g. minimal supersymmetric SM (MSSM), little Higgs, technicolor and two Higgs doublet model (2HDM) [3], [5], [4], [6]. Especially, 2HDM can be an effective theory of the models defined at higher energy scale. The purposes of this paper are

1. the determination of the theory in Higgs sector when Higgs boson(s) is (are) discovered in LHC, and
2. the determination of the current structure in Type-III 2HDM.

If only one neutral Higgs boson is discovered, is that means that the Higgs sector is the SM? The same situation occurs in MSSM and 2HDM if other four Higgs bosons are decoupled since very heavy or weakly coupled. Even if we can determine that the Higgs sector is 2HDM-like by LHC experiment, this model has many currents and couplings, and each coupling constant is a complex parameter of the model. It is very important to determine the absolute values of coupling constants, and relative phases between them.

The lepton flavor violation (LFV) process gives them an answer. These models beyond the SM predict the large LFV [7], [8], [9]. The coupling constant between Higgs boson and fermions tends to be larger proportional to the fermion mass. Especially, the Type-III 2HDM has tree-level flavor changing neutral currents (FCNCs). Moreover, KEK B-factory generates huge number of $\tau^+\tau^-$ pairs. These facts suggest LFV τ decay may appear in near future. In this paper, we study $\tau \rightarrow 3\mu$ mode. The reasons are as follows:

- the Higgs boson can contribute in tree level,
- we expect the clear experimental result since the final state has no photon and no missing particle and
- we can study the polarization information using the initial and final energy momentum distributions [10],[11].

The general Lagrangian for $\tau^+ \rightarrow \mu^+ \mu^+ \mu^-$ decay is written as [10], [11]:

$$\begin{aligned} \mathcal{L} = & -2\sqrt{2}G_F \left\{ g_1(\bar{\tau}_R \mu_L)(\bar{\mu}_R \mu_L) + g_2(\bar{\tau}_L \mu_R)(\bar{\mu}_L \mu_R) \right. \\ & + g_3(\bar{\tau}_R \gamma_\alpha \mu_R)(\bar{\mu}_R \gamma^\alpha \mu_R) + g_4(\bar{\tau}_L \gamma_\alpha \mu_L)(\bar{\mu}_L \gamma^\alpha \mu_L) \\ & \left. + g_5(\bar{\tau}_R \gamma_\alpha \mu_R)(\bar{\mu}_L \gamma^\alpha \mu_L) + g_6(\bar{\tau}_L \gamma_\alpha \mu_L)(\bar{\mu}_R \gamma^\alpha \mu_R) \right\}, \quad (1) \\ & - 2\sqrt{2}G_F m_\tau \left\{ A_R \bar{\tau}_R \sigma^{\alpha\beta} \mu_L F_{\alpha\beta} + A_L \bar{\tau}_L \sigma^{\alpha\beta} \mu_R F_{\alpha\beta} \right\} \\ & + \bar{\mu}(iD^\alpha \gamma_\alpha - m_\mu)\mu - \frac{1}{4}F^{\alpha\beta}F_{\alpha\beta}, \end{aligned}$$

where m_μ and m_τ are the masses of the $\mu^\pm$ and $\tau^\pm$, respectively; G_F is the Fermi constant; $\{\bar{\tau}_L, \bar{\mu}_L, \mu_R\}$ and $\{\bar{\tau}_R, \bar{\mu}_R, \mu_L\}$ are the Dirac spinors $\{\bar{\tau}, \bar{\mu}, \mu\}$ with the helicity operators, $(1 \pm \gamma_5)/2$, respectively; $\sigma^{\alpha\beta} = \frac{i}{2}(\gamma^\alpha \gamma^\beta - \gamma^\beta \gamma^\alpha)$; $D^\alpha = \partial^\alpha + ieA^\alpha$; $F^{\alpha\beta} = \partial^\alpha A^\beta - \partial^\beta A^\alpha$; A^α is the photon field; $e = -|e|$ is the electron charge; A_L and A_R are the complex coefficients of interactions in which the intermediate photon has the left polarization and the right polarization, respectively; and $g_1, \dots, g_6$ are the complex coefficients of various 4 Fermi type interactions.

According to Ref. [11], we can determine the observables,

$$\begin{aligned} a_\pm &= \frac{|g_1|^2}{16} \pm \frac{|g_2|^2}{16} + |g_3|^2 \pm |g_4|^2 \\ b_\pm &= |g_5|^2 \pm |g_6|^2 \\ c_\pm &= |eA_R|^2 \pm |eA_L|^2 \\ d_\pm &= -(Re[g_3 e A_L^*] \pm Re[g_4 e A_R^*]) \\ e_\pm &= -(Re[g_6 e A_R^*] \pm Re[g_5 e A_L^*]) \\ f_+ &= -(Im[g_3 e A_L^*] + Im[g_4 e A_R^*]) \\ g_+ &= -(Im[g_6 e A_R^*] + Im[g_5 e A_L^*]). \end{aligned} \quad (2)$$

a_+ , b_+ , c_+ , d_+ and e_+ are determined from the final-state muon energy distribution and a_- , b_- , c_- , d_- , e_- , f_+ and g_+ are determined from the final-state angular distribution. So, our first task is to determine $g_1, \dots, g_6$, A_L and A_R in this model.

This paper is organized as follows. In Section 2, we derive the coupling constants $g_1, \dots, g_6$, A_L and A_R in this model. In Section 3, we give four scenarios and study the features of each scenario. In Section 4, we discuss the difference between the minimal supersymmetric SM (MSSM) and Type-III 2HDM. In Section 5, we the summary and discussion.

2 Effective Coupling Constants in the Model

In Type-III 2HDM, $\tau \rightarrow 3\mu$ decay can be written in tree level. However, we also consider the one-loop radiative diagrams since the resonance effect enhances the

contribution as follows. These diagrams contain the photon propagator, which is proportional to $1/q^2$, where q is the propagating momentum. The minimum value of q^2 is $4m_\mu^2$. This is realized when the pair created muon anti-muon have the same momentum. On the other hand, in the Higgs mediated diagrams, this part is replaced by the Higgs mass squared.

2.1 Four Fermi Diagrams




Figure 1: The tree-level diagram, which corresponds to the coupling constant g_1 .




Figure 2: The tree-level diagram, which corresponds to the coupling constant g_2 .




Figure 3: The tree-level diagram, which corresponds to the coupling constant $2g_5$.




Figure 4: The tree-level diagram, which corresponds to the coupling constant $2g_6$.

Here, we calculate the tree-level diagrams which contain the neutral Higgs bosons in the intermediate state and are written as Figs. 1, 2, 3 and 4. Comparing these diagrams and the general Lagrangian (1), the effective coupling constants g_1, g_2, g_5 and g_6 contain

$$\begin{aligned}
 g_1 &\supset \frac{1}{-2\sqrt{2}G_F} \left(J_{23}^{H^0*} J_{22}^{H^0} \frac{1}{M_{H^0}^2} + J_{23}^{h^0*} J_{22}^{h^0} \frac{1}{M_{h^0}^2} - J_{23}^{A^0*} J_{22}^{A^0} \frac{1}{M_{A^0}^2} \right) \\
 g_2 &\supset \frac{1}{-2\sqrt{2}G_F} \left(J_{32}^{H^0} J_{22}^{H^0} \frac{1}{M_{H^0}^2} + J_{32}^{h^0} J_{22}^{h^0} \frac{1}{M_{h^0}^2} - J_{32}^{A^0} J_{22}^{A^0} \frac{1}{M_{A^0}^2} \right) \\
 g_5 &\supset \frac{1}{-4\sqrt{2}G_F} \left(J_{23}^{H^0*} J_{22}^{H^0} \frac{1}{M_{H^0}^2} + J_{23}^{h^0*} J_{22}^{h^0} \frac{1}{M_{h^0}^2} + J_{23}^{A^0*} J_{22}^{A^0} \frac{1}{M_{A^0}^2} \right) \\
 g_6 &\supset \frac{1}{-4\sqrt{2}G_F} \left(J_{32}^{H^0} J_{22}^{H^0*} \frac{1}{M_{H^0}^2} + J_{32}^{h^0} J_{22}^{h^0*} \frac{1}{M_{h^0}^2} + J_{32}^{A^0} J_{22}^{A^0*} \frac{1}{M_{A^0}^2} \right),
 \end{aligned} \tag{3}$$

where

$$\begin{aligned}
J_{ij}^{H^0} &= -\frac{gm_i\delta_{ij}}{2\sin\beta M_W}\sin\alpha + \frac{\eta_{ij}^E}{\sqrt{2}\sin\beta}\sin(\alpha-\beta) \\
J_{ij}^{h^0} &= -\frac{gm_i\delta_{ij}}{2\sin\beta M_W}\cos\alpha + \frac{\eta_{ij}^E}{\sqrt{2}\sin\beta}\cos(\alpha-\beta) \\
J_{ij}^{A^0} &= \frac{gm_i\delta_{ij}}{2\sin\beta M_W}\cos\beta + \frac{\eta_{ij}^E}{\sqrt{2}\sin\beta}
\end{aligned} \tag{4}$$

are the effective coupling constants, where g is the $SU(2)_L$ gauge coupling constant; m_i are the i -th family charged lepton masses; α is mixing angle between neutral CP even Higgses; β is defined as $\tan\beta = v_2/v_1$, where v_1 and v_2 are the vacuum expectation values of down and up-sector Higgses, respectively; M_W is the weak boson mass; H^0 and h^0 are the heavier and lighter CP even neutral Higgses, respectively; A^0 is the CP odd neutral Higgs; M_{H^0} , M_{h^0} and M_{A^0} are H^0 , h^0 and A^0 masses, respectively; η_{ij}^E are non-diagonal leptonic Yukawa couplings of Type-III 2HDM. Here, we note that g_5 and g_6 are generated via Fierz transformation. They are the same as g_1 and g_2 in their FCNC parts and flipping the helicity in their flavor conserving neutral current part, respectively.

2.2 Radiative diagrams

We consider the radiative one-loop diagrams since the resonance effect enhances its contribution as mentioned in the first of this section.

In general, the anti-fermion to anti-fermion and imaginary photon FCNC amplitude which momenta are p , $p-q$ and q , respectively, is written as

$$\bar{v}(p)i\left[q^2\gamma^\mu(C_1P_R + C_2P_L) + q^\mu(C_3P_R + C_4P_L) + i\sigma^{\nu\mu}q_\nu(C_5P_R + C_6P_L)\right]v(p-q), \tag{5}$$

where $C_1, \dots, C_6$ are the complex coefficients which should be given by the model; $P_R, P_L \equiv (1 \pm \gamma^5)/2$; $\bar{v}(p)$ and $v(p-q)$ are the spinors of initial and final anti-fermion, respectively.

Considering $\tau \rightarrow 3\mu$ decay, $q^\mu(C_3P_R + C_4P_L)$ term will vanish when we multiply the electromagnetic current $-ie\bar{\mu}\gamma_\mu\mu$ and use the Dirac equation. By the same prescription, $q^2\gamma^\mu(C_1P_R + C_2P_L)$ term becomes

$$\bar{v}(p)i\left[q^2\gamma^\mu(C_1P_R + C_2P_L)\right]v(p-q)\frac{-ig_{\mu\nu}}{q^2}\bar{u}(p_3)(-ie\gamma^\nu)(P_R + P_L)v(p_2), \tag{6}$$

where p_2 and p_3 are the final-state fermion and anti-fermion momenta, respectively. So, $g_3, \dots, g_6$ contain C_1 and C_2 as

$$\begin{aligned}
g_3 &\supset \frac{eC_1}{2\sqrt{2}G_F}, \quad g_4 \supset \frac{eC_2}{2\sqrt{2}G_F} \\
g_5 &\supset \frac{eC_1}{2\sqrt{2}G_F}, \quad g_6 \supset \frac{eC_2}{2\sqrt{2}G_F}.
\end{aligned} \tag{7}$$




Figure 5: The radiative one-loop diagram which have charged Higgs bosons and neutrinos in the loop.




Figure 6: The radiative one-loop diagram which have neutral Higgs bosons and charged leptons in the loop. The dashed line represents H^0 , h^0 and A^0 propagator.

Also, $i\sigma^{\nu\mu}q_\nu(C_5P_R + C_6P_L)$ should be compared with A_L and A_R . They contain C_5 and C_6 as

$$A_L \supset \frac{C_5}{-4\sqrt{2}G_F m_\tau}, \quad A_R \supset \frac{C_6}{-4\sqrt{2}G_F m_\tau}. \quad (8)$$

In Type-III 2HDM, these diagrams are written in Figs. 5 and 6. The coefficients C_1, C_2, C_5 and C_6 are calculated in Appendix C, and the explicit form of effective coupling constants $g_1, \dots, g_6$, A_L and A_R are written in Appendix A.

3 Scenarios

Here, we suppose four possible scenarios in type-III 2HDM. They are as follows:

1. All Higgs bosons except for h^0 are very heavy and decoupled.
2. All Higgs bosons except for A^0 are very heavy and decoupled.
3. $H^\pm$ and A^0 are very heavy and decoupled, where $H^\pm$ are the charged Higgs bosons which have ± 1 electromagnetic charges, respectively. h^0 and H^0 have the masses $M_{h^0} = 98\text{GeV}$ and $M_{H^0} = 115\text{GeV}$, respectively. ZZh^0 coupling is too small to be discovered in LEP experiment.
4. All Higgs bosons except for $H^\pm$ are very heavy and decoupled.

Each scenario suggests different final-state distributions. Comparing them to the experimental result, we can test the Type-III 2HDM.

3.1 Scenario 1: only h^0 is light

If we discover only one neutral Higgs boson in LHC, there may be no signal of new physics in LHC. However, if the LFV is discovered, we have to reconsider whether the Higgs sector is the SM one or not.

In this scenario, we set $M_{H^0}, M_{A^0}, M_{H^\pm} \gg M_{h^0}$, where $M_{H^\pm}$ is $H^\pm$ mass. This situation becomes important when we discover only one neutral Higgs boson in LHC.

According to the above mass relation and Appendix A, we find the relations

$$g_5 \simeq \frac{g_1}{2} \frac{J_{22}^{h^0}}{J_{22}^{h^0*}} + g_3, \quad g_6 \simeq \frac{g_2}{2} \frac{J_{22}^{h^0*}}{J_{22}^{h^0}} + g_4. \quad (9)$$

Substituting Eq. (2) for g_3, g_4, g_5 and g_6 in these relations, and then solving about $|g_1|^2$ and $|g_2|^2$, respectively, they are written by the observables as follows:

$$\begin{aligned} |g_1|^2 &\simeq \frac{8}{25} \left(5(a_+ + a_-) + 3(b_+ + b_-) - 2(e_+ - e_-) \frac{5(d_+ + d_-) - (e_+ - e_-)}{c_+ - c_-} \right) \\ &\quad \pm \frac{16}{25} \sqrt{(b_+ + b_-) - \frac{(e_+ - e_-)^2}{c_+ - c_-}} \sqrt{4(5(a_+ + a_-) - (b_+ + b_-)) - \frac{(5(d_+ + d_-) - (e_+ - e_-))^2}{c_+ - c_-}}, \\ |g_2|^2 &\simeq \frac{8}{25} \left(5(a_+ - a_-) + 3(b_+ - b_-) - 2(e_+ + e_-) \frac{5(d_+ - d_-) - (e_+ + e_-)}{c_+ + c_-} \right) \\ &\quad \pm \frac{16}{25} \sqrt{(b_+ - b_-) - \frac{(e_+ + e_-)^2}{c_+ + c_-}} \sqrt{4(5(a_+ - a_-) - (b_+ - b_-)) - \frac{(5(d_+ - d_-) - (e_+ + e_-))^2}{c_+ + c_-}}. \end{aligned} \quad (10)$$

These equations have the discrete ambiguities in their later terms. These correspond to the sign of imaginary part of $g_5 e A_L^*$ and $g_6 e A_R^*$, respectively. Moreover, using these quantities, we determine

$$|g_3|^2 \simeq \frac{a_+ + a_-}{2} - \frac{|g_1|^2}{16}, \quad |g_4|^2 \simeq \frac{a_+ - a_-}{2} - \frac{|g_2|^2}{16}, \quad (11)$$

and

$$\begin{aligned} \text{Arg}[g_3 e A_L^*] &\simeq \arccos \left[-\frac{d_+ + d_-}{\sqrt{2}|g_3|\sqrt{c_+ - c_-}} \right] \\ \text{Arg}[g_4 e A_R^*] &\simeq \arccos \left[-\frac{d_+ - d_-}{\sqrt{2}|g_4|\sqrt{c_+ + c_-}} \right] \\ \text{Arg}[g_1 \frac{J_{22}^{h^0}}{J_{22}^{h^0*}} g_3^*] &\simeq \arccos \left[\frac{8(b_+ + b_-) - 8(a_+ + a_-) - 3|g_1|^2}{16|g_1||g_3|} \right] \\ \text{Arg}[g_2 \frac{J_{22}^{h^0*}}{J_{22}^{h^0}} g_4^*] &\simeq \arccos \left[\frac{8(b_+ - b_-) - (a_+ - a_-) - 3|g_2|^2}{16|g_2||g_4|} \right] \\ \text{Arg}[g_3 g_5^*] &\simeq \arccos \left[\frac{(b_+ + b_-) + 10|g_3|^2 - 4(a_+ + a_-)}{2\sqrt{2}|g_3|\sqrt{b_+ + b_-}} \right] \\ \text{Arg}[g_4 g_6^*] &\simeq \arccos \left[\frac{(b_+ - b_-) + 10|g_4|^2 - 4(a_+ - a_-)}{2\sqrt{2}|g_4|\sqrt{b_+ - b_-}} \right]. \end{aligned} \quad (12)$$

3.1.1 Situation division

This scenario contains three types of couplings as follows:

1. $J_{31}^{h^0} J_{12}^{h^0}, J_{31}^{h^0} J_{21}^{h^0*}, J_{13}^{h^0} J_{12}^{h^0}, J_{13}^{h^0} J_{12}^{h^0*},$
2. $J_{32}^{h^0} J_{22}^{h^0}, J_{32}^{h^0} J_{22}^{h^0*}, J_{23}^{h^0} J_{22}^{h^0}, J_{23}^{h^0} J_{22}^{h^0*},$
3. $J_{33}^{h^0} J_{32}^{h^0}, J_{33}^{h^0} J_{23}^{h^0*}, J_{33}^{h^0} J_{32}^{h^0}, J_{33}^{h^0} J_{23}^{h^0*}.$

These are divided by the family of intermediate leptons. The tree-level contribution is contained in the second-type coupling. However, we cannot neglect other contributions since each $J_{ij}^{h^0}$ is only a parameter of the model.

To determine which contributions are dominant, we first divide in four cases using the observables which are defined in Eq. (2) and the relations (9) as

$$\frac{a_+}{b_+} \simeq \frac{|g_1|^2 + |g_2|^2 + 16(|g_3|^2 + |g_4|^2)}{4(|g_1|^2 + |g_2|^2) + 16(|g_3|^2 + |g_4|^2) + 16\text{Re}[g_1 \frac{J_{22}^{h^0}}{J_{22}^{h^0*}} g_3^* + g_2 \frac{J_{22}^{h^0*}}{J_{22}^{h^0}} g_4^*]}, \quad (13)$$

d_+/c_+ and c_+/a_+ .

case 1; $a_+/b_+ \simeq 1/4$

When $a_+/b_+ \simeq 1/4$, $|g_1|^2 + |g_2|^2$ is dominant and this means that the tree-level contribution i.e. second-type coupling works. So, even if all $|J_{ij}^{h^0}|$ are the same order, the observable a_+/b_+ suggests this case. We note here that $c_+/a_+ \ll 1$ in this case.

case 2; $a_+/b_+ \simeq 1$ and $c_+/a_+ = \mathcal{O}(1)$

When $a_+/b_+ \simeq 1$, then $|g_3|^2 + |g_4|^2 \gg |g_1|^2 + |g_2|^2$. In this case, the second-type coupling $|J_{23}^{h^0} J_{22}^{h^0*}|^2 + |J_{32}^{h^0} J_{22}^{h^0}|^2$ is highly suppressed compared with other two types of couplings. From the equations in Appendix A, $|J_{23}^{h^0} J_{22}^{h^0*}|^2 + |J_{32}^{h^0} J_{22}^{h^0}|^2$ is highly suppressed as

$$\mathcal{O}(10^{-5}) \gg \frac{|J_{23}^{h^0} J_{22}^{h^0*}|^2 + |J_{32}^{h^0} J_{22}^{h^0}|^2}{|J_{13}^{h^0} J_{12}^{h^0}|^2 + |J_{31}^{h^0} J_{21}^{h^0*}|^2 + |J_{33}^{h^0} J_{32}^{h^0}|^2 + |J_{33}^{h^0} J_{23}^{h^0*}|^2}. \quad (14)$$

The left hand side in above relation is small since the tree-level diagrams affect $|g_1|^2 + |g_2|^2$, while only one-loop diagrams affect $|g_3|^2 + |g_4|^2$. The radiative one-loop currents are dominant to the tree-level FCNC. So, $\tau \rightarrow \mu\gamma$ may be discovered earlier than $\tau \rightarrow 3\mu$. However, $\tau \rightarrow 3\mu$ mode has the advantage that we can determine which of the third-type couplings and the first-type couplings are dominant using the observable c_+/a_+ . If the third-type couplings are dominant, we have an observable relation from Appendix A as,

$$\frac{c_+}{a_+} \simeq \frac{9}{4} + \frac{27}{8} \left(1 - \frac{\text{Re}[J_{33}^{h^0 2}]}{|J_{33}^{h^0}|^2} \right) \frac{3 + 2 \log[\frac{m_\tau^2}{M_{h^0}^2}]}{(4 + 3 \log[\frac{m_\tau^2}{M_{h^0}^2}])^2}. \quad (15)$$

When we set $M_{h^0} = 115\text{GeV}$,

$$\frac{9}{4} - 0.21 \lesssim \frac{c_+}{a_+} \lesssim \frac{9}{4}. \quad (16)$$

In this case, there is one more simple relation,

$$\frac{d_+}{e_+} \simeq 1. \quad (17)$$

We can confirm $|g_3|^2 + |g_4|^2$ dominance using these relations.

case 3; $a_+/b_+ \simeq 1$ **and** $c_+/a_+ = \mathcal{O}(10^{-4})$

If $a_+/b_+ \simeq 1$ as the case 2, however the first-type couplings are dominant and no tuning,

$$\frac{c_+}{a_+} \simeq \frac{9}{16} \left(-1 + 3 \log \left[\frac{q^2}{M_{h^0}^2} \right] \right)^{-2} \simeq 0.00034, \quad (18)$$

where $M_{h^0} = 115\text{GeV}$. These two cases have different order of c_+/a_+ values. This fact is an advantage to determine the current structure.

In this case, we note

$$\frac{d_+}{e_+} \simeq 1 \quad (19)$$

as in case 2.

case 4; $a_+/b_+ \not\simeq 1, 1/4$

If $a_+/b_+ \not\simeq 1$ or $1/4$, this means $|g_1|^2 + |g_2|^2 \sim |g_3|^2 + |g_4|^2$. This case is similar to the case 2 and the case 3 with no tuning, and

$$\mathcal{O}(10^{-5}) \sim \frac{|J_{23}^{h^0*} J_{22}^{h^0*}|^2 + |J_{32}^{h^0} J_{22}^{h^0}|^2}{|J_{13}^{h^0*} J_{12}^{h^0}|^2 + |J_{31}^{h^0} J_{21}^{h^0*}|^2 + |J_{33}^{h^0*} J_{32}^{h^0}|^2 + |J_{33}^{h^0} J_{23}^{h^0*}|^2}. \quad (20)$$

3.1.2 Bounds from Total Branching Ratio

We give the upper limits on the effective couplings in each case studied above. Supposing parity and CP conservation for simplicity, the total branching ratio is written as [11]

$$\text{Br}(\tau^+ \rightarrow \mu_1 \mu_2 \mu_3) = \text{Br}(\tau \rightarrow \mu \nu \bar{\nu}) \left[2a_+ + b_+ + 16d_+ + 8e_+ + \frac{8}{3} \left(24 \log \left[\frac{3}{4\delta} \right] - 13 \right) c_+ \right], \quad (21)$$

where $\delta = 2m_\mu/m_\tau$.




Figure 7: Relation between α and β in scenario 1.

To reduce the free parameter of the model, we consider the relation between α and β . In this scenario, according to the Appendix B, we set $M_{11}M_{22} = M_{12}^2$ since $M_{H^0} \gg M_{h^0}$. Here, we set λ_6 is large enough since $M_{A^0}^2 = \lambda_6(v_1^2 + v_2^2)$ is decoupled, and for CP conservation of Higgs potential, $\lambda_6 = \lambda_5$. So, $\lambda_6 = \lambda_5 \gg \lambda_1, \dots, \lambda_3$ and

$$\begin{aligned} M_{11} &= v_2^2 \lambda_5 \\ M_{22} &= v_1^2 \lambda_5 \\ M_{12} &= M_{21} = \lambda_5 v_1 v_2. \end{aligned} \tag{22}$$

Here, we set the α region as follows:

$$M_{A^0}^2 > 0 \Rightarrow \lambda_6 = \lambda_5 > 0 \Rightarrow M_{12} > 0 \Rightarrow \sin 2\alpha > 0 \Rightarrow 0 < \alpha < \frac{\pi}{2}. \tag{23}$$

Using these conditions, we have the relations from Appendix B as,

$$\begin{aligned} \tan \alpha &= \sqrt{\frac{M_{22}}{M_{11}}}. \\ \frac{1}{\tan^2 \alpha} &= \frac{M_{11}}{M_{22}} \simeq \frac{v_2^2}{v_1^2} = \tan^2 \beta, \\ \cos \alpha &= \sin \beta \\ \cos(\alpha - \beta) &= 2 \sin \beta \cos \beta. \end{aligned} \tag{24}$$

This is explained in Fig. 7

Second-type coupling: $J_{23}^{h^0} J_{22}^{h^0}$

We here consider the case that second-type coupling i.e. $J_{23}^{h^0} J_{22}^{h^0}$ is dominant, which case corresponds to the case 1 in above. In this case, only $g_1 = g_2 =$

$2g_5 = 2g_6$ are considerable and the branching ratio is given as

$$\text{Br}(\tau^+ \rightarrow \mu_1 \mu_2 \mu_3) \simeq 4.8 \times (\eta_{23}^E \eta_{22}^E)^2 \cos^4 \beta \left(\frac{100 \text{GeV}}{M_{h^0}} \right)^4. \quad (25)$$

If we set $M_{h^0} = 115 \text{GeV}$, $\cos \beta = 1/\sqrt{2}$ and $\text{Br}(\tau \rightarrow 3\mu) < 3.2 \times 10^{-8}$ from [12], then the upper limit is written as

$$|\eta_{23}^E \eta_{22}^E| < 0.00022. \quad (26)$$

First-type coupling: $J_{13}^{h^0} J_{12}^{h^0}$

We here consider the case that the first-type coupling i.e. $J_{13}^{h^0} J_{12}^{h^0}$ is dominant, which case corresponds to the case 3 in above. Then $g_1, g_2 \simeq 0$, $g_5 = g_6 = g_3 = g_4$, $eA_L = eA_R$ and the branching ratio is written as

$$\begin{aligned} & \text{Br}(\tau^+ \rightarrow \mu_1 \mu_2 \mu_3) \\ & \simeq 0.14 \times 10^{-5} \times \left[0.73 + (-12.3 + \log[(\frac{100 \text{GeV}}{M_{h^0}})^2])^2 \right] \left(\frac{100 \text{GeV}}{M_{h^0}} \right)^4 \cos^4 \beta (\eta_{13}^E \eta_{12}^E)^2. \end{aligned} \quad (27)$$

If we set $M_{h^0} = 115 \text{GeV}$, $\cos \beta = 1/\sqrt{2}$ and $\text{Br}(\tau \rightarrow 3\mu) < 3.2 \times 10^{-8}$, we give an upper limit as

$$|\eta_{13}^E \eta_{12}^E| \leq 0.032. \quad (28)$$

Third-type coupling: $J_{33}^{h^0} J_{32}^{h^0}$

We here consider the case that the third-type coupling $J_{33}^{h^0} J_{32}^{h^0}$ is dominant, which case corresponds to the case 2 in above. Then $g_1, g_2 \simeq 0$, $g_5 = g_6 = g_3$ and

$$\begin{aligned} & \text{Br}(\tau^+ \rightarrow \mu_1 \mu_2 \mu_3) \\ & \simeq 1.1 \times 10^{-4} (\cos^2 \beta (-\frac{0.0051}{\cos \beta} + \eta_{33}^E \eta_{23}^E)^2 (6.7 - \log[(\frac{100 \text{GeV}}{M_{h^0}})^2])^2 \left(\frac{100 \text{GeV}}{M_{h^0}} \right)^4. \end{aligned} \quad (29)$$

If we set $M_{h^0} = 115 \text{GeV}$, $\cos \beta = 1/\sqrt{2}$ and $\text{Br}(\tau \rightarrow 3\mu) < 3.2 \times 10^{-8}$, we give an upper limit as

$$|(-0.0072 + \eta_{33}^E \eta_{23}^E)| \leq 0.0065. \quad (30)$$

3.2 Scenario 2: only A^0 is light

Here, we study the scenario 2, in which all Higgs bosons except for A^0 are decoupled. In this case, g_5 and g_6 are written from Appendix A as

$$g_5 \simeq -\frac{g_1}{2} \frac{J_{22}^{A^0}}{J_{22}^{A^0*}} + g_3, \quad g_6 \simeq -\frac{g_2}{2} \frac{J_{22}^{A^0*}}{J_{22}^{A^0}} + g_4. \quad (31)$$

The observable difference from the scenario 1 presents in a_+/c_+ when $J_{33}^{A^0} J_{32}^{A^0}$ is dominant and CP violation in the Lagrangian is small. When $J_{13}^{A^0} J_{12}^{A^0}$ or its hermit conjugate (h.c.) is dominant, the difference from $J_{13}^{h^0} J_{12}^{h^0}$ or its h.c. dominant situation is suppressed by m_e/m_τ . When $J_{23}^{A^0} J_{22}^{A^0}$ or its h.c. is dominant, the difference from $J_{23}^{h^0} J_{22}^{h^0}$ or its h.c. dominant situation is suppressed by the fine-structure constant $\alpha_{QED} = e^2/(4\pi)$.

In scenario 1, only h^0 is light, if CP symmetry of Lagrangian is conserved, $Re[(J_{33}^{h^0})^2] = |J_{33}^{h^0}|^2$ and

$$\frac{c_+}{a_+} \simeq \frac{9}{4}, \quad (32)$$

from Eq. (15). On the other hand, in this scenario, only A^0 is light, if CP symmetry of Lagrangian is conserved,

$$\frac{c_+}{a_+} \simeq \frac{9}{4} + \frac{27}{8} \left(1 + \frac{Re[J_{33}^{A^0}]^2}{|J_{33}^{A^0}|^2} \right) \frac{3 + 2 \log[\frac{m_\tau^2}{M_{A^0}^2}]}{(4 + 3 \log[\frac{m_\tau^2}{M_{A^0}^2}])^2} \simeq \frac{9}{4} - 0.21, \quad (33)$$

where we set $M_{A^0} = 115\text{GeV}$. This value is stable even if M_{A^0} is varied intensely since it is in logarithmic function.

3.3 Scenario 3: only H^0 is seen

In this scenario, we can detect only H^0 . We suppose that h^0 , which is lighter than H^0 , could not be detected in LEP since the ZZh^0 coupling in 2HDM is too weak. $H^\pm$ and A^0 are also decoupled since they are very heavy. The LEP experiment suggests a possibility of $M_{H^0} = 115\text{GeV}$ and $M_{h^0} = 98\text{GeV}$ [13]. In this scenario, we set from Appendix B as,

$$\begin{aligned} \frac{M_{11}}{M_{22}} &\simeq \tan^2 \beta \\ \lambda_5 = \lambda_6 &\gg \lambda_{1,2,3} \\ \cos 2\alpha &\simeq -\frac{M_{H^0} + M_{h^0}}{M_{H^0} - M_{h^0}} \cos 2\beta = -\frac{213}{17} \cos 2\beta. \end{aligned} \quad (34)$$

This relation between α and β suggests that $\beta \simeq \pi/4$ in all α range. So, the effective coupling constant $J_{ij}^{H^0}$ is written as

$$J_{ij}^{H^0} \simeq -\frac{gm_i \delta_{ij}}{\sqrt{2}M_W} \sin \alpha + \eta_{ij}^E \sin(\alpha - \frac{\pi}{4}). \quad (35)$$

Also, from Refs. [13] and [14], we set $|g_{ZZh^0}^{2HDM}| < |g_{ZZH}^{SM}|/2$, where $g_{ZZh^0}^{2HDM}$ and g_{ZZH}^{SM} are the ZZh^0 coupling in Type-III 2HDM and ZZ Higgs coupling in the SM, respectively. This suggests the allowed region $|\sin(\beta - \alpha)| < 1/2$ since ZZh^0 term in 2HDM Lagrangian is $M_Z^2 \sin(\beta - \alpha) ZZh^0 / \sqrt{v_1^2 + v_2^2}$, where

M_Z is the Z boson mass. Using the fact $\beta \simeq \pi/4$, the allowed region of α becomes $\frac{\pi}{12} < \alpha < \frac{5\pi}{12}$. After all, the allowed region of α and β in this scenario is explained as in Fig. 8.

Comparing between Fig. 7 and Fig. 8, this scenario has narrower allowed region than the scenario 1. So, if we can determine α or β , we can distinguish this scenario and scenario 1.




Figure 8: The allowed region of α and β in scenario 3. The upper left and the lower right are forbidden by the LEP constraint i.e. $\beta - \pi/6 < \alpha < \beta + \pi/6$. If $M_{h^0} = 98\text{GeV}$ and $M_{H^0} = 115\text{GeV}$, only on the curved line near $\beta = \pi/4$ is allowed.

3.4 Scenario 4: only H^+ is light

We consider all the Higgs bosons except for H^+ are decoupled. Here, we don't consider the parity and CP conservation. In this case, we have the relations from Appendix A as,

$$\begin{aligned} g_1 &\simeq g_2 \simeq g_4 \simeq g_6 \simeq 0 \\ g_3 &\simeq g_5 \simeq \frac{4}{3}eA_L \simeq \frac{4}{3}\frac{m_\tau}{m_\mu}eA_R \simeq -\frac{\alpha_{QED}}{2\sqrt{2}G_F}\frac{(J^{H^+\dagger}J^{H^+})_{32}}{9(4\pi)M_{H^+}^2}, \end{aligned} \quad (36)$$

and the observables become

$$\begin{aligned} c_+ &\simeq -c_- \simeq \frac{9}{16}|g_3|^2 \\ a_\pm &\simeq b_\pm \simeq |g_3|^2 \\ d_\pm &\simeq \pm e_\pm \simeq -\frac{3}{4}|g_3|^2. \end{aligned} \quad (37)$$

$$\begin{aligned}
& \text{Br}(\tau^+ \rightarrow \mu_1 \mu_2 \mu_3) \\
& \simeq 5.3 \times 10^{-8} \frac{1}{\sin^4 \beta} \left(\frac{100 \text{GeV}}{M_{H^+}} \right)^2 |(\eta_{13}^{E*} \eta_{12}^E + \eta_{23}^{E*} \eta_{22}^E + (-0.01 \cos \beta + \eta_{33}^{E*}) \eta_{32}^E)|^2
\end{aligned} \tag{38}$$

If we set $M_{H^\pm} = 100 \text{GeV}$, $\cos \beta = \sin \beta = 1/\sqrt{2}$ and $\text{Br}(\tau \rightarrow 3\mu) < 3.2 \times 10^{-8}$, we give an upper limit as

$$|(\eta_{13}^{E*} \eta_{12}^E + \eta_{23}^{E*} \eta_{22}^E + (-0.0072 + \eta_{33}^{E*}) \eta_{32}^E)| < 0.39 \tag{39}$$

4 Difference Between SUSY and Type-III 2HDM

4.1 MSSM

In most MSSM models, the dominant contribution is from the radiative diagrams [10], [15]. So, at first, the LFV event will be discovered in radiative mode e.g. $\tau \rightarrow \mu \gamma$.

According to section 2.2, this feature appears in the effective coupling constants as

$$g_1 = g_2 = 0, \quad g_3 = g_5, \quad g_4 = g_6, \tag{40}$$

in $\tau \rightarrow 3\mu$ mode. So, the observables $a_\pm$, $b_\pm$, $d_\pm$, $e_\pm$, f_+ and g_+ have the relations,

$$a_\pm = b_\pm, \quad d_\pm = \pm e_\pm, \quad f_+ = g_+ = 0. \tag{41}$$

We note here that imaginary part becomes zero since g_3 , g_4 , eA_L and eA_R have the same complex phases.

On the other hand, the tree-level diagram contributions in type-III 2HDM derives the characteristic relations, $g_1 \neq 0$ and $g_2 \neq 0$. Furthermore, in generally, tree-level heavy gauge boson in intermediate state derives the relation, $a_\pm \neq b_\pm$. In these case we can easily discriminate the difference between MSSM and 2HDM.

Even if the experimental result suggests the relation (41), it may still be Type-III 2HDM. In this case, another observable a_+/c_+ becomes important to distinguish them. If there is no tuning, this situation is realized when $J_{22} = 0$ or $J_{23}^* = J_{32} = 0$. So, if $a_+/c_+ \sim 9/4$ or ~ 0.0034 as derived in section 3.1, 2HDM is strongly supported. On the other hand, if a_+/c_+ has other values, then the MSSM is supported.

4.2 Babu-Kolda Model

In Babu-Kolda model (BKM) [8], MSSM neutral higgses propagate the intermediate state of the $\tau \rightarrow 3\mu$ decay. If we consider the situation, $Br(\tau \rightarrow$

$\mu\gamma) \lesssim Br(\tau \rightarrow 3\mu)$, the radiative diagrams affect little in $\tau \rightarrow 3\mu$ decay. So, we now neglect it. In this model, g_1 and g_5 are written as

$$\begin{aligned} g_1 &= \frac{m_\tau m_\mu \kappa_{32}}{2 \cos^3 \beta} \left(\frac{\cos(\alpha - \beta) \sin \alpha}{M_h^2} + \frac{\sin(\alpha - \beta) \cos \alpha}{M_H^2} - \frac{\sin \beta}{M_A^2} \right) \\ g_5 &= \frac{1}{2} \frac{m_\tau m_\mu \kappa_{32}}{2 \cos^3 \beta} \left(\frac{\cos(\alpha - \beta) \sin \alpha}{M_h^2} + \frac{\sin(\alpha - \beta) \cos \alpha}{M_H^2} + \frac{\sin \beta}{M_A^2} \right), \end{aligned} \quad (42)$$

and others are zero. So, at first, we should check whether or not

$$a_+ = a_-, \quad b_+ = b_-, \quad c_\pm = d_\pm = e_\pm = 0. \quad (43)$$

If h^0 or A^0 is decoupled, we also have to check whether or not the relation

$$a_+ = a_- = 4b_+ = 4b_- \quad (44)$$

is satisfied.

Type-III 2HDM can also make the similar situation as BKM. Even if so, we have a procedure to distinguish them when both of h^0 and A^0 are not decoupled. In Type-III 2HDM, from the relation,

$$\frac{1}{M_{H^0}^2} < \left(\frac{\sin^2(\alpha - \beta)}{M_{H^0}^2} + \frac{\cos^2(\alpha - \beta)}{M_{h^0}^2} \right) < \frac{1}{M_{h^0}^2}, \quad (45)$$

the observable

$$\frac{b_+ - 4a_+}{b_+ + 4a_+} = \frac{2 \left(\frac{\sin^2(\alpha - \beta)}{M_{H^0}^2} + \frac{\cos^2(\alpha - \beta)}{M_{h^0}^2} \right) \frac{1}{M_{A^0}^2}}{\left(\frac{\sin^2(\alpha - \beta)}{M_{H^0}^2} + \frac{\cos^2(\alpha - \beta)}{M_{h^0}^2} \right)^2 + \left(\frac{1}{M_{A^0}^2} \right)^2} \quad (46)$$

is constrained. Similarly, in Babu model, the relation

$$\begin{aligned} -\frac{1}{M_{H^0}^2} &\leq \left(\frac{\cos(\alpha - \beta) \sin \alpha}{M_h^2} + \frac{\sin(\alpha - \beta) \cos \alpha}{M_H^2} \right) \frac{1}{\sin \beta} \\ &= \frac{(M_{H^0}^2 + M_{h^0}^2) \frac{\sin(2\alpha - \beta)}{\sin \beta} + (M_{H^0}^2 - M_{h^0}^2)}{2M_{H^0}^2 M_{h^0}^2} \leq \infty \end{aligned} \quad (47)$$

constrains the observable

$$\frac{b_+ - 4a_+}{b_+ + 4a_+} = \frac{2 \left(\frac{\cos(\alpha - \beta) \sin \alpha}{M_h^2} + \frac{\sin(\alpha - \beta) \cos \alpha}{M_H^2} \right) \frac{1}{\sin \beta} \frac{1}{M_A^2}}{\left(\frac{\cos(\alpha - \beta) \sin \alpha}{M_h^2} + \frac{\sin(\alpha - \beta) \cos \alpha}{M_H^2} \right)^2 \left(\frac{1}{\sin \beta} \right)^2 + \left(\frac{1}{M_A^2} \right)^2}. \quad (48)$$

Figs. 9, 10 and 11 are the allowed regions of this observable in BKM and Type-III 2HDM. In these figures, the yellow region is the allowed region in the BKM, and the blue region is the allowed region in the Type-III 2HDM. Fig. 9 suggests the allowed regions in the condition, $M_{h^0} = 98\text{GeV}$ and $M_{H^0} = 115\text{GeV}$. Similarly, Fig. 10 suggests the allowed regions in the condition, $M_{h^0} = 120\text{GeV}$ and $M_{H^0} = 150\text{GeV}$; and Fig. 11 suggests the allowed regions in the condition, $M_{h^0} = 120\text{GeV}$ and $M_{H^0} = 1000\text{GeV}$. In each case, the allowed region in the Type-III 2HDM is smaller than that of the BKM.




Figure 9: $M_{h^0} = 98\text{GeV}$, $M_{H^0} = 115\text{GeV}$, $98 \leq M_{A^0} \leq 200\text{GeV}$. The yellow region is the allowed region in the Babu model, and the blue region is the allowed region in the Type-III 2HDM.

5 Summary and Discussion

We studied the energy distributions and angular distributions of $\tau \rightarrow 3\mu$ decay products supposing the type-III 2HDM. We supposed four scenarios in section 3. Each scenario has the different feature which we can check using the experimental observables e.g. a_+ , b_+ , a_+/b_+ , c_+/a_+ ,... We explained the upper limits on the effective coupling constants in some cases of the scenario 1. We suggested that the difference between the MSSM case and the type-III 2HDM case can be observed in energy and angular distributions. Also, we suggested that the difference between the BKM case and the type-III 2HDM case can be checked by the observable $(b_+ - 4a_+)/(b_+ + 4a_+)$.

$\tau \rightarrow 3\mu$ decay is a pure leptonic LFV event. Pure leptonic event has no QCD ambiguity and LFV is a null test. So, the experimental results become clear and sensitive to the new physics.

The discovery of LFV event is very important, itself. What we want to know next is the source of it. This paper focus the type-III 2HDM as a source. Comparing it to the MSSM and BKM, we suggest that the energy and angular distribution are the effective probe to distinguish the models.

Even if no new particle except for one Higgs boson is discovered in LHC, the $\tau \rightarrow 3\mu$ events and its energy and angular distributions support the new physics in Higgs sector. High energy collider like LHC is the primary approach to the new physics study. Our analysis is the complimentary approach to them.

The type-III 2HDM has many source of CP violation. This analysis manifests its effects in the observables. The non-zero values of f_+ or g_+ defined in Eq. (2) or $(b_+ \pm b_-)(c_+ \mp c_-) - (e_+ \mp e_-)^2$ means the CP violation.

As a next to LHC high energy collider, the international linear collider (ILC) is planned in near future [16]. If the source of LFV is the Higgs bosons discovered




Figure 10: $M_{h^0} = 120\text{GeV}$, $M_{H^0} = 150\text{GeV}$, $100 \leq M_{A^0} \leq 200\text{GeV}$. The color condition is the same as that of Fig. 9.

in LHC, the ILC will create a number of LFV events since the Higgs bosons becomes the on-shell particle. However, this analysis enables us to precisely determine the next to SM before the ILC era.

References

- [1] M. Spira, A. Djouadi, D. Graudenz, P.M. Zerwas *Nucl. Phys.* **B 453** 17 (1995), e-Print: hep-ph/9504378.
- [2] H. Hufel, G. Pocsik, *Z. Phys.* **C 8** 13 (1981).
- [3] Stephen P. Martin, e-Print: hep-ph/9709356.
- [4] Edward Farhi, Leonard Susskind, *Phys. Rept.* **74** 277 (1981), Koichi Yamawaki, Masako Bando, Ken-iti Matumoto, *Phys. Rev. Lett.* **56** 1335 (1986).
- [5] Martin Schmaltz, David Tucker-Smith, *Ann. Rev. Nucl. Part. Sci.* **55** 229 (2005), e-Print: hep-ph/0502182.
- [6] For a review see J. Gunion, H. Haber, G. Kane and S. Dawson, *The Higgs Hunter's Guide*, (Addison-Wesley, New York, 1990).
- [7] S.-G. Kim, N. Maekawa, A. Matsuzaki, K. Sakurai, and T. Yoshikawa, *Phys. Rev.* **D 75**, 115008 (2007).
- [8] K.S. Babu, Christopher Kolda, *Phys. Rev. Lett.* **89**, 241802 (2002) preprint hep-ph/0206310.
- [9] Rodolfo A. Diaz, R. Martinez, J. Alexis Rodriguez, *Phys. Rev.* **D 64**, 033004 (2001), e-Print: hep-ph/0103050.




Figure 11: $M_{h^0} = 120\text{GeV}$, $M_{H^0} = 1000\text{GeV}$, $100 \leq M_{A^0} \leq 1500\text{GeV}$. The color condition is the same as that of Fig. 9.

- [10] Yasuhiro Okada, Ken-ichi Okumura, Yasuhiro Shimizu, *Phys. Rev. D* **61**, 094001 (2000), preprint hep-ph/9906446.
- [11] A. Matsuzaki, A.I. Sanda, *Phys. Rev. D* **77**, 073003 (2008), e-Print: arXiv:0711.0792.
- [12] Belle Collaboration, *Phys. Lett. B* **660** 154 (2008), e-Print: arXiv:0711.2189.
- [13] R. Barate et al. (LEP Working Group for Higgs boson searches), *Phys. Lett. B* **565**, 61 (2003).
- [14] S.-G.Kim, N.Maekawa, A.Matsuzaki, K.Sakurai, A.I.Sanda, and T.Yoshikawa, *Phys. Rev. D* **74**, 115016 (2006).
- [15] Andrea Brignole, Anna Rossi, *Nucl. Phys. B* **701** 3 (2004). e-Print: hep-ph/0404211
- [16] Nima Arkani-Hamed, Hsin-Chia Cheng, Jonathan L. Feng, Lawrence J. Hall, *Phys. Rev. Lett.* **77**, 1937 (1996) ; M. Cannoni, S. Kolb, O. Panella, *Phys. Rev. D* **68**, 096002 (2003) ; P.M. Ferreira, R.B. Guedes, R. Santos, *Phys. Rev. D* **75**, 055015 (2007).

A Coupling Constants of the Model

$g_1, \dots, g_6$, A_L and A_R in Type-III 2HDM are written as follows:

$$\begin{aligned}
 g_1 &= \frac{1}{-2\sqrt{2}G_F} \left(J_{23}^{H^0*} J_{22}^{H^0} \frac{1}{M_{H^0}^2} + J_{23}^{h^0*} J_{22}^{h^0} \frac{1}{M_{h^0}^2} - J_{23}^{A^0*} J_{22}^{A^0} \frac{1}{M_{A^0}^2} \right), \\
 g_2 &= \frac{1}{-2\sqrt{2}G_F} \left(J_{32}^{H^0} J_{22}^{H^0} \frac{1}{M_{H^0}^2} + J_{32}^{h^0} J_{22}^{h^0} \frac{1}{M_{h^0}^2} - J_{32}^{A^0} J_{22}^{A^0} \frac{1}{M_{A^0}^2} \right),
 \end{aligned} \tag{49}$$

$$\begin{aligned}
g_3 &= \frac{e^2}{2\sqrt{2}G_F} \left[\frac{(J_{13}^{H^0*} J_{12}^{H^0} + J_{23}^{H^0*} J_{22}^{H^0})(-1 + 3\log[\frac{q^2}{M_{H^0}^2}]) + J_{33}^{H^0*} J_{32}^{H^0} (4 + 3\log[\frac{m_\tau^2}{M_{H^0}^2}])}{9(4\pi)^2 M_{H^0}^2} \right. \\
&\quad \left. + (H^0 \rightarrow h^0) + (H^0 \rightarrow A^0) - \frac{(J^{H^+ \dagger} J^{H^+})_{32}}{9(4\pi)^2 M_{H^+}^2} \right], \\
g_4 &= \frac{e^2}{2\sqrt{2}G_F} \left[\frac{(J_{31}^{H^0} J_{21}^{H^0*} + J_{32}^{H^0} J_{22}^{H^0*})(-1 + 3\log[\frac{q^2}{M_{H^0}^2}]) + (J_{33}^{H^0} J_{23}^{H^0*})(4 + 3\log[\frac{m_\tau^2}{M_{H^0}^2}])}{9(4\pi)^2 M_{H^0}^2} \right. \\
&\quad \left. + (H^0 \rightarrow h^0) + (H^0 \rightarrow A^0) \right],
\end{aligned} \tag{50}$$


$$\begin{aligned}
g_5 &= \frac{1}{-4\sqrt{2}G_F} \left(J_{23}^{H^0*} J_{22}^{H^0} \frac{1}{M_{H^0}^2} + (H^0 \rightarrow h^0) + (H^0 \rightarrow A^0) \right) + g_3, \\
g_6 &= \frac{1}{-4\sqrt{2}G_F} \left(J_{32}^{H^0} J_{22}^{H^0*} \frac{1}{M_{H^0}^2} + (H^0 \rightarrow h^0) + (H^0 \rightarrow A^0) \right) + g_4,
\end{aligned} \tag{51}$$

$$\begin{aligned}
A_L &= \frac{e}{-4\sqrt{2}G_F m_\tau} \left[\frac{(J^{H^+ \dagger} J^{H^+})_{32} m_\tau}{6(4\pi)^2 M_{H^+}^2} - \frac{(J^{H^0 \dagger} J^{H^0})_{32} m_\tau}{6(4\pi)^2 M_{H^0}^2} - \frac{(J^{H^0} J^{H^0 \dagger})_{32} m_\mu}{6(4\pi)^2 M_{H^0}^2} \right. \\
&\quad + (H^0 \rightarrow h^0) + (H^0 \rightarrow A^0) \\
&\quad + \frac{(J_{31}^{H^0} J_{12}^{H^0} m_e + J_{32}^{H^0} J_{22}^{H^0} m_\mu)}{2(4\pi)^2 M_{H^0}^2} (-1 + 2\log[\frac{q^2}{M_{H^0}^2}]) \\
&\quad \left. + \frac{J_{33}^{H^0} J_{32}^{H^0} m_\tau}{2(4\pi)^2 M_{H^0}^2} (3 + 2\log[\frac{m_\tau^2}{M_{H^0}^2}]) + (H^0 \rightarrow h^0) - (H^0 \rightarrow A^0) \right], \\
A_R &= \frac{e}{-4\sqrt{2}G_F m_\tau} \left[\frac{(J^{H^+ \dagger} J^{H^+})_{32} m_\mu}{6(4\pi)^2 M_{H^+}^2} - \frac{(J^{H^0 \dagger} J^{H^0})_{32} m_\mu}{6(4\pi)^2 M_{H^0}^2} - \frac{(J^{H^0} J^{H^0 \dagger})_{32} m_\tau}{6(4\pi)^2 M_{H^0}^2} \right. \\
&\quad + (H^0 \rightarrow h^0) + (H^0 \rightarrow A^0) \\
&\quad + \frac{(J_{13}^{H^0*} J_{21}^{H^0*} m_e + J_{23}^{H^0*} J_{22}^{H^0*} m_\mu)}{(4\pi)^2} \frac{-1 + 2\log[\frac{q^2}{M_{H^0}^2}]}{2M_{H^0}^2} \\
&\quad \left. + \frac{J_{33}^{H^0*} J_{23}^{H^0*} m_\tau}{(4\pi)^2} \frac{3 + 2\log[\frac{m_\tau^2}{M_{H^0}^2}]}{2M_{H^0}^2} + (H^0 \rightarrow h^0) - (H^0 \rightarrow A^0) \right],
\end{aligned} \tag{52}$$


where we set $q^2 = 16m_\tau^2 \delta^2 / 9 \simeq 2m_\mu^2$ by the condition

$$x_1 < 1 - \left(\frac{4\delta}{3}\right)^2. \tag{53}$$


Vertex Feynman rules for Type-III 2HDM are as follows:



$$\begin{aligned}
 & \bar{\nu}_{iL} \\
 & \quad \quad \quad H^+ \\
 & \quad \quad \quad \equiv \sqrt{2}iJ_{ij}^{H^+} P_R \\
 & \quad \quad \quad = -\frac{igm_i\delta_{ij}}{\sqrt{2}\sin\beta M_W} \cos\beta P_R + \frac{i\eta_{ij}^E}{\sin\beta} P_R,
 \end{aligned} \tag{54}$$



$$\begin{aligned}
 & E_{jR} \\
 & \bar{E}_{jR} \\
 & \quad \quad \quad H^- \\
 & \quad \quad \quad \equiv \sqrt{2}iJ_{ij}^{H^+*} P_L \\
 & \quad \quad \quad = -\frac{igm_i\delta_{ij}}{\sqrt{2}\sin\beta M_W} \cos\beta P_L + \frac{i\eta_{ij}^{E*}}{\sin\beta} P_L,
 \end{aligned} \tag{55}$$



$$\begin{aligned}
 & \bar{\ell}_{iL} \\
 & \quad \quad \quad \{H^0, h^0, A^0\} \\
 & \quad \quad \quad \equiv iJ_{ij}^{H^0} P_R = -\frac{igm_i\delta_{ij}}{2\sin\beta M_W} \sin\alpha P_R + \frac{i\eta_{ij}^E}{\sqrt{2}\sin\beta} \sin(\alpha - \beta) P_R \\
 & \quad \quad \quad , iJ_{ij}^{h^0} P_R = -\frac{igm_i\delta_{ij}}{2\sin\beta M_W} \cos\alpha P_R + \frac{i\eta_{ij}^E}{\sqrt{2}\sin\beta} \cos(\alpha - \beta) P_R \\
 & \quad \quad \quad , J_{ij}^{A^0} P_R = \frac{gm_i\delta_{ij}}{2\sin\beta M_W} \cos\beta P_R + \frac{\eta_{ij}^E}{\sqrt{2}\sin\beta} P_R,
 \end{aligned} \tag{56}$$

$$\begin{aligned}
& \equiv iJ_{ij}^{H^0*}P_L = -\frac{igm_i\delta_{ij}}{2\sin\beta M_W}\sin\alpha P_L + \frac{i\eta_{ij}^E*}{\sqrt{2}\sin\beta}\sin(\alpha-\beta)P_L \\
& , iJ_{ij}^{h^0*}P_L = -\frac{igm_i\delta_{ij}}{2\sin\beta M_W}\cos\alpha P_L + \frac{i\eta_{ij}^E*}{\sqrt{2}\sin\beta}\cos(\alpha-\beta)P_L \\
& , -J_{ij}^{A^0*}P_L = -\frac{gm_i\delta_{ij}}{2\sin\beta M_W}\cos\beta P_L - \frac{\eta_{ij}^E*}{\sqrt{2}\sin\beta}P_L.
\end{aligned}
\tag{57}$$

B Higgs masses

In Type-III 2HDM, the general Higgs potential is written as [6]:

$$\begin{aligned}
V(\phi_1, \phi_2) &= \lambda_1(\phi_1^\dagger\phi_1 - v_1^2)^2 + \lambda_2(\phi_2^\dagger\phi_2 - v_2^2)^2 \\
&+ \lambda_3[(\phi_1^\dagger\phi_1 - v_1^2) + (\phi_2^\dagger\phi_2 - v_2^2)]^2 \\
&+ \lambda_4[(\phi_1^\dagger\phi_1)(\phi_2^\dagger\phi_2) - (\phi_1^\dagger\phi_2)(\phi_2^\dagger\phi_1)] \\
&+ \lambda_5[Re(\phi_1^\dagger\phi_2) - v_1v_2\cos\xi]^2 \\
&+ \lambda_6[Im(\phi_1^\dagger\phi_2) - v_1v_2\sin\xi]^2,
\end{aligned}
\tag{58}$$

, where $\lambda_1, \dots, \lambda_6$ are the real coefficients; ϕ_1 and ϕ_2 are the Higgs fields with the vacume expectation values, v_1 and v_2 , respectively; and ξ is the CP phase. When we set $\xi = 0$ for CP conservation of Higgs potential, the neutral CP even and odd Higgs masses are written as

$$\begin{aligned}
M_{H^0, h^0} &= \frac{1}{2}[M_{11} + M_{22} \pm \sqrt{(M_{11} - M_{22})^2 + 4M_{12}^2}], \\
M_{A^0}^2 &= \lambda_6(v_1^2 + v_2^2),
\end{aligned}
\tag{59}$$

respectively, where

$$\begin{aligned}
M_{11} &= 4v_1^2(\lambda_1 + \lambda_3) + v_2^2\lambda_5, \\
M_{22} &= 4v_2^2(\lambda_2 + \lambda_3) + v_1^2\lambda_5, \\
M_{12} &= M_{21} = (4\lambda_3 + \lambda_5)v_1v_2.
\end{aligned}
\tag{60}$$

We define the mixing angle α as

$$\tan 2\alpha = \frac{2M_{12}}{M_{11} - M_{22}}.
\tag{61}$$

C Radiative one-loop diagrams


We explain the general charged fermion-charged fermion-photon one-loop diagrams which contain scalar fields in loop.


For convenience, we define the projection operators,

$$P_1 = \frac{1 \pm \gamma_5}{2}, \quad P_2 = \frac{1 \mp \gamma_5}{2}. \quad (62)$$

C.1 Neutral Scalar


We set the general charged fermion-charged fermion-neutral scalar and charged fermion-charged fermion-photon vertices as follows:





where ℓ is the charged fermion; $\ell_{ir} = P_r \ell_i$, where $r = 1, 2$; h^0 is the neutral scalar; V_{ik}^r are the complex coupling constant; and γ^μ is the photon; $Q = -1$ for a lepton.

The one-loop diagrams which contain the neutral scalars are written as:



$$= i\mathcal{M}_2,$$

$$= i\mathcal{M}_3.$$

C.1.1 Neutral Higgs Contribution with $iV_{ik}^1 P_1$ and $iV_{kj}^1 P_1$

$$\begin{aligned}
& i\mathcal{M}_1^\mu + i\mathcal{M}_2^\mu + i\mathcal{M}_3^\mu \Big|_{11} \\
&= \frac{-ieQV_{ik}^1 V_{kj}^1 m_k}{(4\pi)^2} \int dx \int dy \bar{v}(p) \frac{1}{\Delta} \left[(y-1)i\sigma^{\nu\mu} q_\nu P_1 \right. \\
&\quad \left. + \frac{(2x+y-1)q^2}{m_i^2 - m_j^2} \gamma^\mu (m_i P_1 + m_j P_2) + (2x+y-1)q^\mu P_1 \right] v(p-q),
\end{aligned} \tag{63}$$

where

$$\Delta = x(x+y-1)q^2 + y(x+y-1)m_i^2 - xym_j^2 + (1-y)m_k^2 + yM^2; \tag{64}$$

$|_{11}$ in the left hand side of Eq. (63) means that we use $iV_{ik}^1 P_1$ and $iV_{kj}^1 P_1$ vertices; m_i , m_j and m_k are the ℓ_i , ℓ_j and ℓ_k masses, respectively; and M is the h^0 mass.


C.1.2 Neutral Higgs Contribution with $iV_{ik}^2 P_2$ and $iV_{kj}^1 P_1$

$$\begin{aligned}
& i\mathcal{M}_1^\mu + i\mathcal{M}_2^\mu + i\mathcal{M}_3^\mu \Big|_{21} \\
&= \frac{-ieQV_{ik}^2 V_{kj}^1}{(4\pi)^2} \int dx dy \frac{1}{\Delta} \bar{v}(p) \left[\right. \\
&\quad q^2 \gamma^\mu \frac{1}{m_j^2 - m_i^2} \left\{ x(2x + y - 2)m_j^2 P_1 - (x + y - 1)(2x + y)m_i^2 P_1 - y(y + 2x - 1)m_i m_j P_2 \right\} \\
&\quad + (x + y - 1)(2x + y)m_i q^\mu P_1 - x(2x + y - 2)m_j q^\mu P_2 \\
&\quad \left. - i\sigma^{\nu\mu} q_\nu \{ x y m_j P_2 - y(x + y - 1)m_i P_1 \} \right] v(p - q),
\end{aligned} \tag{65}$$

where $|_{21}$ in the left hand side means that we use $iV_{ik}^2 P_2$ and $iV_{kj}^1 P_1$ vertices.

C.2 charged scalar

We set the general charged fermion-neutral fermion-charged scalar and charged scalar-charged scalar-photon vertices as follows:



A Feynman diagram showing the decay of a selectron ($\bar{\ell}_{i1}$) into a lepton (ℓ_{i1}) and a photon (γ). The selectron line enters from the left and splits into a lepton line and a photon line. The lepton line continues to the right, and the photon line is represented by a wavy line. The diagram is labeled with $\bar{\ell}_{i1}$ at the top left and ℓ_{i1} at the bottom left. The photon line is labeled with γ at the top right. The diagram is also labeled with H^- and $= iW_{ik}^2 P_2$ at the bottom right.

$$\bar{\nu}_{i2}$$

$$H^+ = iU_{ik}^1 P_1,$$

$\bar{\nu}_{i1}$
 $H^+ = iU_{ik}^2 P_2,$

$$H^- \quad p_- \quad \gamma^\mu \quad = ieQ(p_+ - p_-)^\mu, \quad H^+ \quad p_+$$

where $H^\pm$ are the charged scalars which have ± 1 electromagnetic charges, respectively; ν_k are the neutral fermions. and $p_\pm$ are the $H^\pm$ momenta, respectively.

The one-loop diagrams which contain the charged scalars are

$$\bar{\ell}_i^+(p) \quad \nu_k(p-k) \quad \ell_j^+(p-q) \quad = i\mathcal{M}'_1, \quad H(k) \quad H(k-q) \quad \gamma_\mu^*(q)$$

$$= i\mathcal{M}'_2,$$

$$= i\mathcal{M}'_3.$$

C.2.1 Charged Higgs Contribution with $iW_{ik}^1 P_1$ and $iU_{kj}^1 P_1$

$$\begin{aligned}
& i\mathcal{M}'_1{}^\mu + i\mathcal{M}'_2{}^\mu + i\mathcal{M}'_3{}^\mu \Big|_{11} \\
&= \frac{-ieQW_{ik}^1 U_{kj}^1 m_k}{(4\pi)^2} \int dx \int dy \bar{v}(p) \frac{1}{\Delta'} \left[yi\sigma^{\nu\mu} q_\nu P_1 \right. \\
&\quad \left. + \frac{(2x+y-1)q^2}{m_i^2 - m_j^2} \gamma^\mu (m_i P_1 + m_j P_2) + (2x+y-1)q^\mu P_1 \right] v(p-q),
\end{aligned} \tag{66}$$

where

$$\Delta' = x(x+y-1)q^2 + y(x+y-1)m_i^2 - xym_j^2 + (1-y)M^2 + ym_k^2; \tag{67}$$

M is the $H^\pm$ mass; and $|_{11}$ in the left hand side of Eq. (66) means that we use $iW_{ik}^1 P_1$ and $iU_{kj}^1 P_1$ vertices.

C.2.2 Charged Higgs Contribution with $iW_{ik}^2 P_2$ and $iU_{kj}^1 P_1$

$$\begin{aligned}
& i\mathcal{M}'_1{}^\mu + i\mathcal{M}'_2{}^\mu + i\mathcal{M}'_3{}^\mu \Big|_{21} \\
&= \frac{-ieQW_{ik}^2 U_{kj}^1}{(4\pi)^2} \int dx dy \frac{1}{\Delta'} \bar{v}(p) \left[\right. \\
&\quad - q^2 \gamma^\mu \frac{2x+y-1}{m_j^2 - m_i^2} \left\{ x(m_j^2 P_1 + m_i m_j P_2) - (x+y-1)(m_i^2 P_1 + m_i m_j P_2) \right\} \\
&\quad - (x+y-1)(2x+y-1)m_i q^\mu P_1 + x(2x+y-1)m_j q^\mu P_2 \\
&\quad \left. + i\sigma^{\nu\mu} q_\nu \left\{ x y m_j P_2 - y(x+y-1)m_i P_1 \right\} \right] v(p-q),
\end{aligned} \tag{68}$$

where $|_{21}$ in the left hand side means that we use $iW_{ik}^2 P_2$ and $iU_{kj}^1 P_1$ vertices.

C.3 Application to the Type-III 2HDM

For the type-III 2HDM radiative one-loop diagrams in which neutral Higgses propagate, we sum over four vertex substitutions written as:

$$V_{ik}^1 P_1 \rightarrow J_{3k}^{\{H^0, h^0, A^0\}} P_R, \quad V_{kj}^1 P_1 \rightarrow J_{k2}^{\{H^0, h^0, A^0\}} P_R, \tag{69}$$

$$V_{ik}^1 P_1 \rightarrow J_{k3}^{\{H^0, h^0, A^0\}*} P_L, \quad V_{kj}^1 P_1 \rightarrow J_{2k}^{\{H^0, h^0, A^0\}*} P_L, \tag{70}$$

$$V_{ik}^2 P_2 \rightarrow J_{k3}^{\{H^0, h^0, A^0\}*} P_L, \quad V_{kj}^1 P_1 \rightarrow J_{k2}^{\{H^0, h^0, A^0\}} P_R, \tag{71}$$

$$V_{ik}^2 P_2 \rightarrow J_{3k}^{\{H^0, h^0, A^0\}} P_R, \quad V_{kj}^1 P_1 \rightarrow J_{2k}^{\{H^0, h^0, A^0\}*} P_L. \tag{72}$$

Also, we set the masses as

$$\begin{aligned}
m_i &\rightarrow m_\tau, \\
m_j &\rightarrow m_\tau, m_\mu, m_e, \\
m_k &\rightarrow m_\mu, \\
M &\rightarrow M_{H^0}, M_{h^0}, M_{A^0},
\end{aligned} \tag{73}$$

and take summation over the subscript j and propagating Higgses H^0, h^0, A^0 .

Similarly, for the diagrams in which charged Higgses propagate, we substitute for the vertices as

$$\begin{aligned} W_{ik}^2 P_2 &\rightarrow \sqrt{2} J_{k3}^{H^+*} P_L, \\ U_{kj}^1 P_1 &\rightarrow \sqrt{2} J_{k2}^{H^+} P_R. \end{aligned} \tag{74}$$

Also, we set the masses as

$$\begin{aligned} m_i &\rightarrow m_\tau, \\ m_j &\rightarrow m_\tau, m_\mu, m_e, \\ m_k &\rightarrow m_\mu, \\ M &\rightarrow M_{H^\pm}, \end{aligned} \tag{75}$$

and take summation over the subscript j .